

## Elab Fluor® 647 Anti-Mouse Perforin Antibody[S16009A]

Catalog Number: GFH1294M

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

|                         |                                                                                                                                                                        |
|-------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reactivity              | Mouse                                                                                                                                                                  |
| Host                    | Rat                                                                                                                                                                    |
| Isotype                 | Rat IgG2a, κ                                                                                                                                                           |
| Clone No.               | S16009A                                                                                                                                                                |
| Isotype Control         | Elab Fluor® 647 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832M]                                                                                               |
| Conjugation             | Elab Fluor® 647                                                                                                                                                        |
| Conjugation Information | Elab Fluor® 647 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter). |
| Storage Buffer          | Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.                                                                                                    |


### Applications

### Recommended usage

#### FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



C57BL/6 murine splenocytes are stained with FITC Anti-Mouse CD49b Antibody and Elab Fluor® 647 Anti-Mouse Perforin Antibody (Left). Splenocytes are stained with FITC Anti-Mouse CD49b Antibody and Elab Fluor® 647 Rat IgG2a, κ Isotype Control (Right).

### Preparation & Storage

|          |                                                                                                                                                           |
|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Storage  | Keep as concentrated solution.<br>This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| Shipping | Ice bag                                                                                                                                                   |

### Antigen Information

|                 |                                                            |
|-----------------|------------------------------------------------------------|
| Alternate Names | PfnPfn-1Pfp;Cytolysin;Lymphocyte pore-forming protein;Prf1 |
| Uniprot ID      | P10820                                                     |
| Gene ID         | 18646                                                      |

### For Research Use Only

## Background

Perforin is a 70 kD cytolytic protein that is expressed in the cytoplasmic granules of cytotoxic T lymphocytes (CTLs) and natural killer (NK) cells. Perforin is one of the major effector molecules used by cytotoxic T cells and NK cells to mediate targeted cell lysis.