A Reliable Research Partner in Life Science and Medicine

Recombinant Human BVES/POPDC1 Protein (GST Tag)

Catalog Number: PKSH030823

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source E.coli-derived Human BVES/POPDC1 protein Met 1-Asn36, with an N-terminal GST

Calculated MW 30.8 kDa
Observed MW 32.2 kDa
Accession NP 671488.1

Bio-activity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin Please contact us for more information.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Ferritin, light polypeptide (FTL) is the light subunit of the ferritin protein. Ferritin is the major intracellular iron storage protein in prokaryotes and eukaryotes. It is composed of 24 subunits of the heavy and light ferritin chains. Storage of iron in the tissues occurs in the form of ferritin and hemosiderin. The latter originates from ferritin that has undergone intracellular digestion of its protein shell, leaving the iron core. Ferritin and hemosiderin are components of a continuum. Ferritin has been identified in all types of living organisms: animals, plants, molds, and bacteria. Whithin the protein shell of ferritin, iron is first oxidized to the ferric state for storage as ferric oxyhdroxide. Thus, ferritin removes excess iron from the cell sap where it could otherwise participate in peroxidation mechanisms.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017