

A Reliable Research Partner in Life Science and Medicine

Recombinant Human IL-17A Protein(Halo Tag)

Catalog Number: PDMH100459

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human IL-17A protein Gly24-Ala155, with an C-terminal Halo

Calculated MW Observed MW 45-50 kDa Q16552 Accession

Not validated for activity **Bio-activity**

Properties

> 95% as determined by reducing SDS-PAGE. **Purity**

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -Storage

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


Shipping This product is provided as lyophilized powder which is shipped with ice packs. **Formulation**

Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human IL-17A proteins,2µg/lane of Recombinant Human IL-17A proteins, was resolved with SDS-PAGE under reducing conditions, showing bands at 45-50 KD

Background

For Research Use Only

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

IL17, also known as IL17a, is a cytokine that belongs to the IL-17 family. Cytokines are proteinaceous signaling compounds that are major mediators of the immune response. They control many different cellular functions including proliferation, differentiation, and cell survival/apoptosis but are also involved in several pathophysiological processes including viral infections and autoimmune diseases. Cytokines are synthesized under various stimuli by a variety of cells of both the innate(monocytes, macrophages, dendritic cells) and adaptive(T- and B-cells) immune systems. The IL-17 family of cytokines includes six members, IL-17/IL-17A, IL-17B, IL-17C, IL-17D, IL-17E/IL-25, and IL-17F, which are produced by multiple cell types. IL-17 regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2(PTGS2/COX-2), as well as enhance the production of nitric oxide(NO). High levels of IL-17 are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis, and multiple sclerosis.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com