



A Reliable Research Partner in Life Science and Medicine

RAMP2 Polyclonal Antibody

catalog number: E-AB-15828

Note: Centrifuge before opening to ensure complete recovery of vial contents.

T		4.0	
Desc	т	atiar	
	9.01	JULUI	

Reactivity Human

Immunogen Synthetic peptide of human RAMP2

Host Rabbit Isotype IgG

PurificationAffinity purificationConjugationUnconjugated

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

WB 1:500-1:2000

Data

100 55 — 40 — 35 — 25 — 15 —

Western Blot analysis of 231 cell using RAMP2 Polyclonal

Antibody at dilution of 1:650

Calculated-MV:20 kDa

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

The protein encoded by this gene is a member of the RAMP family of single-transmembrane-domain proteins, called receptor (calcitonin) activity modifying proteins (RAMPs). RAMPs are type I transmembrane proteins with an extracellular N terminus and a cytoplasmic C terminus. RAMPs are required to transport calcitonin-receptor-like receptor (CRLR) to the plasma membrane. CRLR, a receptor with seven transmembrane domains, can function as either a calcitonin-gene-related peptide (CGRP) receptor or an adrenomedullin receptor, depending on which members of the RAMP family are expressed. In the presence of this (RAMP2) protein, CRLR functions as an adrenomedullin receptor. The RAMP2 protein is involved in core glycosylation and transportation of adrenomedullin receptor to the cell surface.

For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email:techsupport@elabscience.cn