

Recombinant Human CD160 protein(His tag)

Catalog Number:PDMH100041



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|------------------------------------|-----------------------------------|
| Synonyms | Natural killer cell receptor BY55 |
| Species | Human |
| Expression Host | HEK293 Cells |
| Sequence | Met1-Leu158 |
| Accession | O95971 |
| Calculated Molecular Weight | 14.9 kDa |
| Observed molecular weight | 25 kDa |
| Tag | C-His |

Properties

| | |
|-----------------------|---|
| Purity | > 90 % as determined by reducing SDS-PAGE. |
| Endotoxin | Please contact us for more information. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 90 % as determined by reducing SDS-PAGE.

Background

CD16 antigen, also known as Natural killer cell receptor BY55 and CD16, is a cell membrane protein which contains one Ig-like V-type (immunoglobulin-like) domain. CD16 is a GPI-anchored lymphocyte surface receptor in which expression is mostly restricted to the highly cytotoxic CD56(dim)CD16(+) peripheral blood NK subset. CD16 is a receptor showing broad specificity for both classical and non-classical MHC class I molecules. CD16 is expressed in spleen, peripheral blood, and small intestine. Expression of CD16 is restricted to functional NK and T cytotoxic lymphocytes. CD16 acts as a co-activator receptor for CD3-induced proliferation of CD4+ CD16+ T cells isolated from inflammatory skin lesions. Unique CD4+ CD16+ lymphocyte subset may play a role in the pathogenesis of skin inflammation. Activated NK lymphocytes release a soluble form of CD16 that functionally impairs the MHC-I-specific cytotoxic CD8(+) T

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CD160 protein(His tag)

Catalog Number:PDMH100041



lymphocyte responsiveness.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com