


Recombinant Glypican-3 Monoclonal Antibody

catalog number: AN301865L

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human
Immunogen	Recombinant human Glypican-3 fragment
Host	Rabbit
Isotype	Rabbit IgG
Clone	C311487
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.
Applications	Recommended Dilution
WB	1:2000-1:10000
IHC	1:100-1:500
IF	1:50

Data




Western Blot with Glypican-3 Monoclonal Antibody at dilution of 1:10000. Lane 1: Raji (-), Lane 2: HepG2


Observed-MW: 15, 70 kDa
Calculated-MW: 70, 40 kDa



Immunohistochemistry of paraffin-embedded Human liver (Negative tissue) using Glypican-3 Monoclonal Antibody at dilution of 1:500.



Immunohistochemistry of paraffin-embedded Human hepatocellular carcinoma using Glypican-3 Monoclonal Antibody at dilution of 1:500.



Immunofluorescent analysis of (100% Ice-cold methanol) fixed HepG2 cells using anti-Glypican-3 Monoclonal Antibody at dilution of 1:50.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3

Immunofluorescent analysis of (100% Ice-cold methanol)
fixed Raji(Negative cell line) cells using anti-Glypican-3
Monoclonal Antibody at dilution of 1:50.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

Glypican 3 belongs to Glypicans (GPCs), a family of glycosylphosphatidylinositol (GPI)-anchored heparan sulphate proteoglycans (HSPGs) that may play a role in the control of cell division and growth regulation. In mammals, there are six GPCs (GPC1 to GPC6), all of which have a similar core-protein size of approx. 60 kDa and the clustering of glycosaminoglycan attachment site near the C-terminus. They are tethered to the cell surface by GPI linkages, which can be cleaved by endogenous phospholipases, thus releasing the protein. Glypican 3 (GPC3) is highly expressed in many tissues during development and plays an important role in the regulation of embryonic growth. Loss-of-function mutations of GPC3 result in the Simpson-Golabi-Behmel overgrowth syndrome (SGBS), and Gpc-3 null mice display developmental overgrowth. In hepatocellular carcinoma (HCC), the overexpression of glypican 3 has been demonstrated to be a reliable diagnostic indicator.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3