

## Recombinant Human ANP32A/PHAP1 Protein (His & GST Tag)

Catalog Number: PKSH030963

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                      |   |
|----------------------|---|
| <b>Species</b>       | Human   |
| <b>Source</b>        | E.coli-derived Human ANP32A/PHAP1 protein Glu 2-Lys 238, with an N-terminal His & GST |
| <b>Calculated MW</b> | 55.4 kDa  |
| <b>Observed MW</b>   | 50 kDa  |
| <b>Accession</b>     | NP_006296.1   |
| <b>Bio-activity</b>  | Not validated for activity  |

### Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 85 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | Please contact us for more information.   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.                     |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from sterile 20mM Tris, 10% glycerol, 1mM DTT, 0.5mM GSH<br>Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual. |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

### Data



> 85 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

acidic leucine-rich nuclear phosphoprotein 32 family member A, also known as acidic nuclear phosphoprotein pp32, Leucine-rich acidic nuclear protein, Mapmodulin, Potent heat-stable protein phosphatase 2A inhibitor I1PP2A, Putative HLA-DR-associated protein I, PHAPI and ANP32A, is a nucleus, cytoplasm and endoplasmic reticulum. ANP32A/ LANP is expressed in all tissues tested. It is highly expressed in kidney and skeletal muscle, moderate levels of expression is in brain, placenta and pancreas. ANP32A/ LANP is weakly expressed in lung. It is found in all regions of the brain examined (amygdala, caudate nucleus, corpus callosum, hippocampus and thalamus), with highest levels in amygdala. ANP32A/ LANP is a component of the SET complex, which also contains SET, APEX1, HMGB2 and NME1. It directly interacts with SET. ANP32A/ LANP also interacts with ATXN1/SCA1. ANP32A/ LANP is implicated in a number of cellular processes, including proliferation, differentiation, caspase-dependent and caspase-independent apoptosis, suppression of transformation (tumor suppressor), inhibition of protein phosphatase 2A, regulation of mRNA trafficking and stability in association with ELAVL1, and inhibition of acetyltransferases as part of the INHAT (inhibitor of histone acetyltransferases) complex. ANP32A/ LANP plays a role in E4F1-mediated transcriptional repression.