

## Recombinant Mouse VCL/Vinculin Protein (His Tag)

**Catalog Number:** PKSM040490

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	HEK293 Cells-derived Mouse VCL/Vinculin protein Met1-Gln1066, with an C-terminal His
<b>Calculated MW</b>	118.2 kDa
<b>Observed MW</b>	118 kDa
<b>Accession</b>	NP_033528.3
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

Vinculin (VCL) is a cytoskeletal protein that is closely related to both cell-matrix interactions and cell-cell junctions. VCL is a membrane-cytoskeletal protein in focal adhesion plaques that is involved in linkage of integrin adhesion molecules to the actin cytoskeleton. The protein contains an acidic N-terminal domain and a basic C-terminal domain separated by a proline-rich middle segment. This protein has multi-ligand properties and has been found to interact with a number of microfilament associated proteins, such as talin, a-actinin, and paxillin, which reportedly bind to either the head or tail domains of vinculin.

### For Research Use Only