# **Elabscience**®

## Porcine IL-1a Antibody Pair Set

Catalog No.E-KAB-0617ApplicationsELISASynonymsIL1A;IL1-A;IL1;IL1F1;Preinterleukin 1 Alpha;Hematopoietin-1;Pro-Interleukin-1-Alpha

#### **Kit components & Storage**

Title	Specifications	Storage
Porcine IL-1α Capture Antibody	1 vial, 100 µ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Porcine IL-1a Detection Antibody	1 vial, 50 μL	Store at -20°C for one year. Avoid
(Biotin)		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0617)	
		Porcine IL-1α Capture Antibody	Porcine IL-1α Detection Antibody (Biotin)
Immunogen	Immunogen	Recombinant Porcine IL-1a protien	Recombinant Porcine IL-1a protien
Information	Swissprot	P18430	
Product details	Reactivity	Porcine	Porcine
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Porcine IL-1a in ELISAs.	

For Research Use Only

# **Elabscience**®

### Applications

Porcine IL-1a Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Porcine IL-1α Capture	
Capture		Antibody	10
			Optical Density
ELISA	1:1000-1:10000	Porcine IL-1a Detection	
Detection		Antibody (Biotin)	0.1
			100 1000 10000 Porcine IL-1α Concentration ( pg/mL )

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### Background

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is a pleiotropic cytokine involved in various immune responses , inflammatory processes , and hematopoiesis. This cytokine is produced by monocytes and macrophages as a proprotein , which is proteolytically processed and released in response to cell injury , and thus induces apoptosis. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. It has been suggested that the polymorphism of these genes is associated with rheumatoid arthritis and Alzheimer's disease.

For Research Use Only