Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

HSP40-4 Polyclonal Antibody

catalog number: E-AB-11313

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen Recombinant protein of human DNAJA1

Host Rabbit **Isotype** IgG


PurificationAffinity purificationConjugationUnconjugated

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


Applications Recommended Dilution

WB 1:500-1:2000 **IHC** 1:100-1:300

Data




Western Blot analysis of HepG2, Raji, A431 and 231 cell using HSP40-4 Polyclonal Antibody at dilution of 1:800



Immunohistochemistry of paraffin-embedded Human colon cancer using HSP40-4 Polyclonal Antibody at dilution of 1:60

Calculated-MV:45 kDa



Immunohistochemistry of paraffin-embedded Human thyroid cancer using HSP40-4 Polyclonal Antibody at dilution of 1:60

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

Web: www.elabscience.cn

temperature recommended.

Background

For Research Use Only

Tel: 400-999-2100

Elabscience®

Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

DnaJ-like proteins interact with HSP 70 molecular chaperones and function to facilitate protein folding and mitochondrial protein import. HSP 40-4, also known as HDJ2, is the human DnaJ homolog that functions as a co-chaperone with a cysteine-rich zinc finger domain. The cellular redox enzyme thioredoxin interacts with HSP 40-4, and oxidation and reduction reversibly regulate HSP 40-4 function in response to the changing redox states of the cell. The zinc finger domain of HSP 40-4 may act as a redox sensor of chaperone-mediated protein-folding machinery, since HSP 40-4 inactivation leads to the oxidation of cysteine thiols and a simultaneous release of coordinated zinc. Loss of the HSP 40-4 protein may be linked to severe defects in spermatogenesis that involve aberrant androgen signaling.

For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email:techsupport@elabscience.cn