

PIK3CA Polyclonal Antibody

catalog number: **D-AB-10260L**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Rat
Immunogen	Recombinant Human PIK3CA Protein expressed by E.coli
Host	Rabbit
Isotype	IgG
Purification	Antigen Affinity Purification
Buffer	PBS with 0.05% Proclin300, 1% protective protein and 50% glycerol, pH7.4

Applications

Recommended Dilution


WB	1:500-1:1000
IHC	1:100-1:200

Data




Western blot with PIK3CA Polyclonal antibody at dilution of 1:1000. lane 1: JurKat whole cell lysate


Observed-MW: 110 kDa
Calculated-MW: 124 kDa



Immunohistochemistry of paraffin-embedded Rat kidney using PIK3CA Polyclonal Antibody at dilution of 1:100



Immunohistochemistry of paraffin-embedded Human breast using PIK3CA Polyclonal Antibody at dilution of 1:100



Immunohistochemistry of paraffin-embedded Rat kidney using PIK3CA Polyclonal Antibody at dilution of 1:100

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Phosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. The protein encoded by this gene represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has been found to be oncogenic and has been implicated in cervical cancers. A pseudogene of this gene has been defined on chromosome 22.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V2.1