A Reliable Research Partner in Life Science and Medicine

Recombinant Human IL17BR/IL17RB Protein (His Tag)

Catalog Number: PKSH030735

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human IL17BR/IL17RB protein Met 1-Gly289, with an C-

terminal His

 Calculated MW
 31.5 kDa

 Observed MW
 41-45 kDa

 Accession
 NP 061195.2

Bio-activity Immobilized human IL17BR-His at 10 μg/ml (100 μl/well) can bind human Fc-IL25,

The EC₅₀ of human Fc-IL25 is 0.1-0.3 μ g/ml.

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Web:www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

IL17RB (Interleukin 17 Receptor B) is a Protein Coding gene. IL17RB is the receptor for IL17E, the only member of the IL17 family promoting Th2 reactions. IL17RB is induced on human macrophages by IL4 and enhanced by TGFbeta. Overexpression of IL17RB is associated with poor prognosis and the short survival of breast cancer patients. IL17RB/IL17B signaling triggers a substantial increase in cell growth, proliferation, and migration through the activation of NF-kappaB as well as the up-regulation of the Bcl-2. IL17RB may be the only gene expressed in CD4+ T cells whose transcript measurement is correlated with the variation in IgE level in asthmatics. Diseases associated with IL17RB include Chronic Mucocutaneous Candidiasis and Seborrheic Infantile Dermatitis.

Fax: 1-832-243-6017