A Reliable Research Partner in Life Science and Medicine

# **Recombinant Human TGFB3 Protein**

Catalog Number: PKSH033140

Note: Centrifuge before opening to ensure complete recovery of vial contents.

-								
D	es	C	ľ٦	n	tı	or	١	

Species Human

Source HEK293 Cells-derived Human TGFB3 protein Ala301-Ser412(Tyr340Phe)

 Mol\_Mass
 12.7 kDa

 Accession
 P10600

**Bio-activity** Measured by its ability to inhibit the IL-4-dependent proliferation of TF-1 mouse T

cells. The ED50 for this effect is 10-80 pg/ml.

# **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 0.01 EU per  $\mu$ g of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs. **Formulation** Lyophilized from a 0.2 μm filtered solution of 50mM Glycine-HCl, 150mM NaCl, pH

2.5.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

# Data



> 95 % as determined by reducing SDS-PAGE.

#### Background

Transforming growth factor beta 3(TGFB3) is a member of a TGF - $\beta$  superfamily which is defined by their structural and functional similarities. TGFB3 is secreted as a complex with LAP. This latent form of TGFB3 becomes active upon cleavage by plasmin, matrix metalloproteases, thrombospondin -1, and a subset of integrins. It binds with high affinity to TGF- $\beta$  RII, a type II serine/threonine kinase receptor. TGFB3 is involved incell differentiation, embryogenesis and development. It is believed to regulate molecules involved in cellular adhesion and extracellular matrix (ECM) formation during the process of palate development. Without TGF- $\beta$ 3, mammals develop a deformity known as a cleft palate.

#### For Research Use Only