# Recombinant Human KLK7 protein (His Tag)

**Catalog Number: PDEH100824**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description

| | |
|---|---|
| **Species** | Human |
| **Mol_Mass** | 25.3 kDa |
| **Accession** | P49862 |
| **Bio-activity** | Not validated for activity |

## Properties

| | |
|---|---|
| **Purity** | > 95% as determined by reducing SDS-PAGE. |
| **Endotoxin** | < 10 EU/mg of the protein as determined by the LAL method |
| **Storage** | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 ℃. Reconstituted protein solution can be stored at 4-8℃ for 2-7 days. Aliquots of reconstituted samples are stable at < -20℃ for 3 months. |
| **Shipping** | This product is provided as lyophilized powder which is shipped with ice packs. |
| **Formulation** | Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol. |
| **Reconstitution** | It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis. |

## Data




SDS-PAGE analysis of Human KLK7 proteins, 2μg/lane of Recombinant Human KLK7 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 31 KD.

## Background

Human Kallikrein 7 is a member of the tissue kallikrein family of extracellular serine proteases that is made up of 15 members. It is predominantly expressed in the skin. A major physiological function of Kallikrein 7 is to regulate the desquamation process (the shedding of corneocytes from the outer layer of the epidermis) through proteolysis of the intercellular adhesive structures between corneocytes. Dysregulation of Kallikrein 7 has been linked to several inflammatory skin diseases including atopic dermatitis, psoriasis, and Netherton syndrome. Studies have shown that Kallikrein 5 is a potential physiological activator for Kallikrein 7. The proform of Kallikrein 7 can be activated by thermolysin.

**For Research Use Only**