

Mouse PCSK9 Antibody Pair Set

Catalog No.	E-KAB-0553	Applications	ELISA
Synonyms	FH3;HCHOLA3;LDLCQ1;NARC-1;NARC1;PC9		

Kit components & Storage

Title	Specifications	Storage
Mouse PCSK9 Capture Antibody	1 vial, 100 µg	Store at -20℃ for one year. Avoid freeze/thaw cycles.
Mouse PCSK9 Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20℃ for one year. Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0553)	
		Mouse PCSK9 Capture Antibody	Mouse PCSK9 Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Mouse PCSK9 protien	Recombinant Mouse PCSK9 protien
	Swissprot	Q80W65	
Product details	Reactivity	Mouse	Mouse
	Host	Rabbit	Rabbit
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50% glycerol; pH 7.5	PBS with 0.04% Proclin 300; 1% protective protein; 50% glycerol; pH 7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Mouse PCSK9 in ELISAs.	

For Research Use Only

Applications

Mouse PCSK9 Sandwich ELISA Assay

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4 µg/mL	Mouse PCSK9 Capture Antibody	
ELISA Detection	1:1000-1:10000	Mouse PCSK9 Detection Antibody (Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Crucial player in the regulation of plasma cholesterol homeostasis. Binds to low-density lipid receptor family members: low density lipoprotein receptor (LDLR) , very low density lipoprotein receptor (VLDLR) , apolipoprotein E receptor (LRP1/APOER) and apolipoprotein receptor 2 (LRP8/APOER2) , and promotes their degradation in intracellular acidic compartments. Acts via a non-proteolytic mechanism to enhance the degradation of the hepatic LDLR through a clathrin LDLRAP1/ARH-mediated pathway. May prevent the recycling of LDLR from endosomes to the cell surface or direct it to lysosomes for degradation. Can induce ubiquitination of LDLR leading to its subsequent degradation. Inhibits intracellular degradation of APOB via the autophagosome/lysosome pathway in a LDLR-independent manner. Involved in the disposal of non-acetylated intermediates of BACE1 in the early secretory pathway. Inhibits epithelial Na⁺ channel (ENaC) - mediated Na⁺ absorption by reducing ENaC surface expression primarily by increasing its proteasomal degradation. Regulates neuronal apoptosis via modulation of LRP8/APOER2 levels and related anti-apoptotic signaling pathways.

For Research Use Only