



A Reliable Research Partner in Life Science and Medicine

Recombinant CD40 Monoclonal Antibody

catalog number: AN301709L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;

Immunogen Recombinant human CD40 fragment

 Host
 Rabbit

 Isotype
 IgG, κ

 Clone
 A417


Purification Protein A purified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

WB 1:1000-1:5000 **IHC** 1:200-1:1000

Data



Western Blot with CD40 Monoclonal Antibody at dilution of

1:5000. Lane 1: Raji Observed-MW:44-48 kDa Calculated-MW:32 kDa Immunohistochemistry of paraffin-embedded Human Hodgkin lymphoma using CD40 Monoclonal Antibody at dilution of 1:1000.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

The CD40 antigen is a 44-48 kDa type I integral membrane protein of the tumor necrosis factor receptor (TNFR) superfamily. This antigen is found on B cell lines, is strongly expressed by interdigitating cells (IDC), basal epithelial cells, and is also present on macrophages, some endothelial cells, and follicular dendritic cells. It is a well-known regulator of T cell immunity including in cancer. CD40-ligand, primarily expressed on activated T cells, interacts with CD40 on APCs resulting in a 'licensed' state.

For Research Use Only