A Reliable Research Partner in Life Science and Medicine

BCAS3 Polyclonal Antibody

catalog number: E-AB-18703

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse

Immunogen Fusion protein of human BCAS3

Host Rabbit
Isotype IgG


Purification Antigen affinity purification

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution


WB 1:500-1:2000 **IHC** 1:30-1:150

Data



Western blot analysis of 293T cell lysates using BCAS3 Polyclonal Antibody at dilution of 1:500

> Observed-MW:Refer to figures Calculated-MW:101 kDa



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using BCAS3 Polyclonal Antibody at dilution of 1:35(×200)



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using BCAS3 Polyclonal Antibody at dilution of 1:35(×200)

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

BCAS3 (breast carcinoma amplified sequence 3), also designated MAAB or GAOB1, is a 913 amino acid protein that is believed to be involved in breast cancer progression. The gene is regulated by ER (estrogen receptor alpha) and expressed in multiple tissues, including malignant human brain lesions. It is overexpressed and amplified in breast cancer cell lines. BCAS3 contains three WD40 repeat regions, a bromodomain, a rare zinc-finger motif, four probable DNA-binding domains, and two kinase-inducible phosphorylation domains. Five variants are produced due to alternative splicing. BCAS3 interacts with histone H3 and PCAF, which is indicative of histone acetyltransferase activity. BCAS3 also exhibits ER transactiviation activity by acting as a coactivator with PELP1 or MTA1. The amplification and translocation between the BCAS3 gene and the BCAS4 gene results in a fusion transcript is overexpressed in MCF-7 cells.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Rev. V1.8