# Recombinant Human 5'-Nucleotidase/5'-NT/CD73 (AAH65937.1, C-Fc)

Catalog Number: PKSH033885



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
- 1	00	cri	m	17	٦m
J			174	, T. U	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

SpeciesHumanMol\_Mass84.6 kDaAccessionAAH65937.1

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin**  $\leq 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ 

**Storage** Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

**Shipping** This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel


packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 120mM NaCl, 4mM CaCl<sub>2</sub>,

20% Glycerol, pH 7.5.

**Reconstitution** Not Applicable

### Data



> 95 % as determined by reducing SDS-PAGE.

## Background

CD73 is an ecto 5'Nucleotidase expressed by most cell types. CD73 hydrolyzes extracellular nucleotides into membrane permeable nucleosides. CD73 is one of several enzymes responsible for the production of extracellular adenosine, a signaling molecule that is involved in responses to inflammation and tissue injury. CD73 is a lymphocyte maturation marker that has functions independent of its catalytic activity. CD73 is also a regulator of leukocyte extravasation, a function that requires its 5'Nucleotidase activity. Defects in NT5E are the cause of calcification of joints and arteries (CAJA). The recombinant CD73 lacking GPI anchor is secreted as a monomer.

## For Research Use Only