

## FITC Anti-Human CD305 Antibody[NKTA255]

Catalog Number: AN00330C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG1
<b>Clone No.</b>	NKTA255
<b>Isotype Control</b>	FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792C]
<b>Conjugation</b>	FITC
<b>Conjugation Information</b>	FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


### Applications

### Recommended usage

#### FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (millie cells in 100 μL staining volume or per 113 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



Staining of normal human peripheral blood cells with APC Anti-Human CD3 Antibody and FITC Anti-Human CD305 Antibody[NKTA255] (left) or FITC Mouse IgG1, κ Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	LAIR-1
<b>Uniprot ID</b>	Q6GTX8
<b>Gene ID</b>	3903

### For Research Use Only

## Background

CD305, also known as LAIR-1 (leukocyte-associated Ig-like receptor-1), is a 40 kD type I transmembrane glycoprotein. LAIR-1 and LAIR-2 are the only members of the LAIR family in the Ig superfamily. LAIR-1 is an inhibitory molecule characterized by ITIMs in the cytoplasmic domain and collagen ligands. CD305 is expressed on NK cells, T cells, B cells, monocytes, dendritic cells, eosinophils, basophils and mast cells. LAIR-1 functions in the inhibition of cell cytotoxicity, cell activation, proliferation and differentiation.