

Recombinant JNK Monoclonal Antibody

catalog number: E-AB-81477

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse;Rat
Immunogen	Recombinant protein of human JNK1
Host	Rabbit
Isotype	Rabbit IgG
Clone	C473263
Purification	Affinity Purified
Buffer	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.05% stabilizer and 0.05% protective protein.

Applications

Recommended Dilution

WB	1:500-1:1000
IF	1:50-1:100


Data



Western blot detection of JNK1/2/3 in C6,3T3 cell lysates using JNK1/2/3 Rabbit mAb(1:1000 diluted). Predicted band size: 48,53kDa. Observed band size: 46,54kDa.

Observed-MW: 46 /54 kDa

Calculated-MW: 48/53 kDa



Immunofluorescence of JNK1/2/3 (green) in HeLa cells using JNK1/2/3 Rabbit mAb at dilution 1:100, and DAPI(blue)

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

MAPK8(Mitogen-activated protein kinase 8) is also named as JNK1,PRKM8,SAPK1,SAPK1C and belongs to the MAP kinase subfamily. MAPK8 is activated by dual phosphorylation at a Thr-Pro-Tyr motif during response to UV light. MAPK8 functions to phosphorylate c-Jun at N-terminal serine regulatory sites of Ser-63 and Ser-73, mapping within the transactivation domain. Phosphorylation of these sites in response to UV results in transcriptional activation of c-Jun. It has 4 isoforms produced by alternative splicing with the molecular weight of 46 kDa and 48 kDa. This protein can be phosphorylated and this antibody recognizes the 46 kDa and 55 kDa bands in western blot.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.9