



A Reliable Research Partner in Life Science and Medicine

HNRNPR Polyclonal Antibody

catalog number: E-AB-92340

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen A synthetic peptide of human HNRNPR


Host Rabbit Isotype IgG

Purification Affinity purification

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Recommended Dilution Applications WB 1:500-1:2000 1:50-1:200 IF 1:50-1:200 IHC

Data



Western blot analysis of extracts of various cell lines using HNRNPR Polyclonal Antibody at 1:1000 dilution.


Observed-MW:Refer to figures Calculated-MW:59 kDa/66 kDa/70 kDa/71 kDa



Immunohistochemistry of paraffin-embedded human esophagus using HNRNPR Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with (40x lens). Perform microwave antigen retrieval with 10 mM 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded rat heart using HNRNPR Polyclonal Antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining



Immunohistochemistry of paraffin-embedded mouse liver using HNRNPR Polyclonal Antibody at dilution of 1:100 PBS buffer pH 7.2 before commencing with IHC staining protocol.


For Research Use Only

Fax: 1-832-243-6017

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine



Immunofluorescence analysis of U2OS cells using HNRNPR Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

This gene encodes an RNA-binding protein that is a member of the spliceosome C complex, which functions in premRNA processing and transport. The encoded protein also promotes transcription at the c-fos gene. Alternative splicing results in multiple transcript variants. There are pseudogenes for this gene on chromosomes 4, 11, and 10.

Fax: 1-832-243-6017