
Recombinant Human LTBR/TNFRSF3 Protein (Fc Tag)

Catalog Number: PKSH032718

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|----------------|--|
| Species | Human |
| Source | HEK293 Cells-derived Human LTBR; TNFRSF3 protein Gln31-Met227, with an C- |
| | terminal Fc |
| Calculated MW | 48.8 kDa |
| Observed MW | 61 kDa |
| Accession | P36941 |
| Bio-activity | Not validated for activity |
| Properties | |
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 |
| | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of |
| | reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. |
| | Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants |
| | before lyophilization. |
| | Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Tumor necrosis factor receptor superfamily member 3; also known as Lymphotoxin-beta receptor; Tumor necrosis factor C receptor; Tumor necrosis factor receptor 2-related protein; Tumor necrosis factor receptor type III; LTBR; TNFCR; TNFR3 and TNFRSF3; is a member of the tumor necrosis factor (TNF) family of receptors. LTBR is a single-pass type I membrane protein and contains four TNFR-Cys repeats. It is expressed on the surface of most cell types; but not on T and B lymphocytes. LTBR and its ligand play a role in the development and organization of lymphoid tissue and transformed cells. Activation of LTBR can trigger apoptosis. In addition; LTBR can lead to the release of the cytokine interleukin 8.

For Research Use Only

Tel:400-999-2100