# **Elabscience**®

### Human IL-5 Antibody Pair Set

Catalog No.	E-KAB-0470	Applications	ELISA
Synonyms	IL5;EDF;TRF;E-CSF;ESGF		

#### **Kit components & Storage**

Title	Specifications	Storage
Human IL-5 Capture Antibody	1 vial, 100 µ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Human IL-5 Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20°C for one year. Avoid
		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0470)		
		Human IL-5 Capture Antibody	Human IL-5 Detection Antibody (Biotin)	
Immunogen	Immunogen	Recombinant Human IL-5 protien	Recombinant Human IL-5 protien	
Information	Swissprot	P05113		
Product details	Reactivity	Human	Human	
	Host	Rabbit	Rabbit	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol; pH 7.5	protective protein; 50% glycerol; pH	
			7.5	
	Purify	Antigen Affinity	Antigen Affinity	
	Specificity	Detects Human IL-5 in ELISAs.		

For Research Use Only

## **Elabscience**®

#### Applications

Human IL-5 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Human IL-5 Capture	
Capture		Antibody	10
			» <sup>1</sup>
			all Demait
ELISA	1:1000-1:10000	Human IL-5 Detection	O 0.1
Detection		Antibody (Biotin)	
			0.01 -
			Human IL-5 Concentration ( pg/mL )

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

#### Background

This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. The encoded cytokine plays a major role in the regulation of eosinophil formation , maturation , recruitment and survival. The increased production of this cytokine may be related to pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receptor , which is a heterodimer , whose beta subunit is shared with the receptors for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene is located on chromosome 5 within a cytokine gene cluster which includes interleukin 4 (IL4) , interleukin 13 (IL13) , and CSF2. This gene , IL4 , and IL13 may be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31.

For Research Use Only