Recombinant Human PFKFB3 (N-6His)

Catalog Number: PKSH033971



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
- 1	00	cri	m	17	٦m
J			174	, T. U	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

 Species
 Human

 Mol_Mass
 61.8 kDa

 Accession
 Q16875

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu\text{g of the protein as determined by the LAL method.}$

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel


packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM PB, 50mM KCl, 5mM DTT, 0.1mM

EDTA, 50% glycerol, 0.1% TritonX-100, pH 6.0.

Reconstitution Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Fructose-2,6-biphosphatase 3, also known as 6-phosphofructo-2-kinase or PFK2 or PFKFB3, involved in both the synthesis and degradation of fructose-2,6-bisphosphate, a regulatory molecule that controls the activity of the enzymes phosphofructokinase 1 (PFK-1) and fructose 1,6-bisphosphatase (FBPase-1) to regulate glycolysis and gluconeogenesis. Highly phosphorylated PFKFB3 protein was found in human tumor cells, vascular endothelial cells, and smooth muscle cells. Because of its proto-oncogenic character, the PFK-2/FBPase-2 of the PFKFB3 gene is assumed to play a critical role in tumorigenesis. PFKFB3 also plays a crucial role in the progression of cancerous cells by enabling their glycolytic pathways even under severe hypoxic conditions, which makes it a potential target for cancer therapy.

For Research Use Only