IL8 Monoclonal Antibody

catalog number: E-AB-22204



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Synthetic C-TERM Peptide of IL-8

HostMouseIsotypeIgGClone13F8

Purification Protein A purification

Conjugation Unconjugated


buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein

protectant and 50% glycerol.

Applications Recommended Dilution

IHC 1:100-200

Data





Immunohistochemistry of paraffin-embedded Human lung carcinoma tissue using IL8 Monoclonal Antibody at dilution of 1:200.

Immunohistochemistry of paraffin-embedded Human kidney tissue using IL8 Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

IL-8 is a chemotactic factor that attracts neutrophils, basophils, and T-cells, but not monocytes. It is also involved in neutrophil activation. It is released from several cell types in response to an inflammatory stimulus. IL-8(6-77) has a 5-10-fold higher activity on neutrophil activation, IL-8(5-77) has increased activity on neutrophil activation and IL-8(7-77) has a higher affinity to receptors CXCR1 and CXCR2 as compared to IL-8(1-77), respectively.

For Research Use Only