

A Reliable Research Partner in Life Science and Medicine

Recombinant CD25 Monoclonal Antibody

catalog number: AN301472L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;

Immunogen Recombinant human CD25 fragment

HostRabbitIsotypeIgG, κCloneA167

Purification Protein A purified


Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.


Applications Recommended Dilution

IF 1:50


FCM 1:50-1:100

Data





Immunofluorescent analysis of (4% Paraformaldehyde) fixed Immunofluorescent analysis of (4% Paraformaldehyde) fixed Jurkat cells using anti-CD25 Monoclonal Antibody at dilution of 1:50.



Flow cytometric analysis of human CD25 expression on Jurkat cells. Cells were stained with purified anti-Human CD25, then a Alexa Fluor 488-conjugated second step antibody. The histogram were derived from events with the forward and side light-scatter characteristics of intact cells.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping lce bag

Background

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Rev. V1.0

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains produce a medium-affinity receptor. Normally an integral-membrane protein, soluble IL2RA has been isolated and determined to result from extracellular proteolyisis. Alternately-spliced IL2RA mRNAs have been isolated, but the significance of each is presently unknown. Mutations in this gene are associated with interleukin 2 receptor alpha deficiency. Patients with severe Coronavirus Disease 2019 (COVID-19), the disease caused by the novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), have significantly elevated levels of IL2R in their plasma. Similarly, serum IL-2R levels are found to be elevated in patients with different types of carcinomas. Certain IL2RA and IL2RB gene polymorphisms have been associated with lung cancer risk.

For Research Use Only