

Recombinant Rhesus Macaque CD47/IAP/OA3 (N-6His-Flag)



Catalog Number:PKSQ050106

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Leukocyte Surface Antigen CD47;Antigenic Surface Determinant Protein OA3;Integrin-Associated Protein;IAP;Protein MER6;CD47;MER6
Species	Rhesus Macaque
Expression Host	HEK293 Cells
Sequence	Gln19-Glu141
Accession	F7A802
Calculated Molecular Weight	15.7 kDa
Observed molecular weight	33-55 kDa
Tag	N-His-Flag

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD47, also known as Integrin α Associated Protein (IAP) and OA3, is a glycosylated atypical member of the immunoglobulin superfamily. Mouse CD47 is an integral membrane protein that consists of a extracellular domain (ECD) with a single Ig γ -like domain, five membrane-spanning regions with short intervening loops, and C α -terminal cytoplasmic tail. CD47 has a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins. It plays an important role in memory formation and synaptic plasticity in the hippocampus. As a receptor for SIRPA, its binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. Interaction with SIRPB1 mediates cell-cell adhesion, it enhances superantigen-dependent T-cell-mediated

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Rhesus Macaque CD47/IAP/OA3 (N-6His-^{dg} Flag)



Catalog Number:PKSQ050106

proliferation and costimulates T-cell activation. It may play a role in membrane transport and/or integrin dependent signal transduction. It also prevents premature elimination of red blood cells.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017