## **Elabscience Biotechnology Co., Ltd.**





# AF/LE Purified Anti-Mouse CD3 Antibody[17A2]

Catalog Number: GF10130

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

**Reactivity** Mouse

Immunogen Recombinant Mouse CD3 protein

**Host** Rat

**Isotype** Rat IgG2b, κ

Clone 17A2

**Purification** >98%, Protein A/G purified

Conjugation None (AF/LE)


Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the

LAL method.

### Applications Recommended Dilution

FCM  $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$ 

#### Data



C57/BL6 Mouse splenocytes were stained with 0.2 µg AF/LE Purified Anti-Mouse CD3 Antibody[17A2] (Right) and 0.2 µg

Rat IgG2b, κ Isotype Control (Left), followed by Elab Fluor<sup>®</sup> 647-conjugated Goat Anti-Rat IgG Secondary Antibody, then anti-Mouse CD8 FITC-conjugated Monoclonal Antibody.

#### **Preparation & Storage**

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid

freeze / thaw cycles. This preparation contains no preservatives, thus it

should be handled under as eptic conditions.

Shipping Ice bag

#### **Background**

CD3, also known as T3, is a member of the lg superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3 is composed of CD3 $\epsilon$ ,  $\delta$ ,  $\gamma$  and  $\zeta$  chains. It forms a TCR complex by associating with TCR  $\alpha/\beta$  or  $\gamma/\delta$  chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and antigen recognition by binding the peptide/MHC antigen complex.