ARIH2 Polyclonal Antibody

catalog number: E-AB-19064



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse

Immunogen Fusion protein of human ARIH2

Host Rabbit Isotype IgG

Purification Antigen affinity purification


Conjugation Unconjugated

buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

WB 1:1000-1:5000 **IHC** 1:50-1:300

Data




Western blot analysis of Human heart tissue lysate using ARIH2 Polyclonal Antibody at dilution of 1:1000

Immunohistochemistry of paraffin-embedded Human tonsil tissue using ARIH2 Polyclonal Antibody at dilution of $1:80(\times 200)$

Observed-MV: Refer to figures

Calculated-MV:58 kDa



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using ARIH2 Polyclonal Antibody at dilution of $1:80(\times 200)$

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack,upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

ARIH2 Polyclonal Antibody

catalog number: E-AB-19064



TRIAD1, also known as ARIH2 (ariadne homolog 2) or ARI2, is a 493 amino acid protein that contains one IBR-type zinc finger and two RING-type zinc fingers and belongs to the ariadne subfamily of RBR proteins. Localized to the nucleus, TRIAD1 interacts with UBE2L3 and is thought to act as an E3 ubiquitin-protein ligase, functioning to accept ubiquitin from E2 ubiquitin-conjugating enzymes and transfer the acquired ubiquitin residue to target substrates. TRIAD1 is subject to post-translational DNA damage-dependent phosphorylation, probably by ATM or ATR. The gene encoding TRIAD1 maps to human chromosome 3, which houses over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of human cancer-related gene loci.