

A Reliable Research Partner in Life Science and Medicine

Recombinant MYH7 Monoclonal Antibody

catalog number: AN301987L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Rat; Mouse

Immunogen Peptide. This information is proprietary to PTMab.

 Host
 Rabbit

 Isotype
 IgG, κ

 Clone
 A707


Purification Protein Apurified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.


Applications Recommended Dilution

IHC 1:1000-1:2000

Data




Immunohistochemistry of paraffin-embedded Human cardiac muscle using MYH7 Monoclonal Antibody at dilution of 1:2000.




Immunohistochemistry of paraffin-embedded Human skeletal muscle using MYH7 Monoclonal Antibody at dilution of 1:2000.



Immunohistochemistry of paraffin-embedded Human colon(Negative tissue) using MYH7 Monoclonal Antibody at dilution of 1:2000.



Immunohistochemistry of paraffin-embedded Mouse skeletal muscle using MYH7 Monoclonal Antibody at dilution of 1:2000.



Immunohistochemistry of paraffin-embedded Rat skeletal muscle using MYH7 Monoclonal Antibody at dilution of 1:2000.

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com



Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Shipping Ice bag

Background

Nonmuscle myosin is an actin-based motor protein essential to cell motility, cell division, migration, adhesion, and polarity. The holoenzyme consists of two identical heavy chains and two sets of light chains. The light chains (MLCs) regulate myosin II activity and stability. The heavy chains (NMHCs) are encoded by three genes, MYH9, MYH10, and MYH14, which generate three different nonmuscle myosin II isoforms, IIa,IIb, and IIc, respectively. While all three isoforms perform the same enzymatic tasks, binding to and contracting actin filaments coupled to ATPhydrolysis, their cellular functions do not appear to be redundant and they have different subcellular distributions. The carboxy-terminal taildomain of myosin II is important in isoform-specific subcellular localization. Research studies have shown that phosphorylation of myosin IIa at Ser1943 contributes to the regulation of breast cancer cell migration.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.1