

APC Anti-Human/Mouse IL-5 Antibody[TRFK5]

Catalog Number: E-AB-F1205UE

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Mouse
Host	Rat
Isotype	Rat IgG1, κ
Clone No.	TRFK5
Isotype Control	APC Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09823E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 $\mu\text{g}/10^6$ cells in 100 μL volume].
------------	---

Data



HEK293T cells transiently transfected with pcDNA3.1 plasmid encoding Mouse IL-5 gene are stained with APC Anti-Human/Mouse IL-5 Antibody (filled gray histogram) or APC Rat IgG1, κ Isotype Control (empty black histogram).

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	B-cell differentiation factor I;IL-5;Interleukin-5;T-cell replacing factor;TRF
Uniprot ID	P05113;P04401
Gene ID	16191;3567

For Research Use Only

Background

IL-5 is a homodimeric, disulphide-linked protein produced by T-cells. Monomeric human IL-5 is a 126 amino acid protein with a reported molecular weight of 26 kD for the homodimeric protein. Mouse and human IL-5 are approximately 70% identical. IL-5 has been shown to promote the growth of immature hematopoietic BFU-E progenitors, stimulate the activation and differentiation of eosinophils, and promote the generation of cytotoxic lymphocytes. Mouse IL-5 induces the differentiation and proliferation of pre-activated B-cells and stimulates the production and secretion of IgM and IgA by B-cells stimulated with bacterial endotoxin. The TRFK5 antibody can neutralize the bioactivity of natural or recombinant IL-5.