## **Elabscience**®


## FITC Anti-Mouse CD106 Antibody[M/K-2.7]

Catalog Number: E-AB-F1091C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
Isotype	Rat lgG1, κ
Clone No.	WK-2.7
Isotype Control	FITC Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09822C]
Conjugation	FITC
Conjugation Information	FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical
	filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 $\mu$ L of antibody per test (million cells in 100 $\mu$ L staining volume or per 100 $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



C57BL/6 murine bone marrow cells are stained with FITC Anti-Mouse CD106 Antibody (filled gray histogram). Unstained bone marrow cells (empty black histogram) are used as control.

Preparation & Storag	e
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged
	exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	CD106;V-CAM1;VCAM-1;Vascular cell adhesion protein 1;Vcam1
Uniprot ID	P29533
Gene ID	22329

## For Research Use Only

## **Elabscience**®

Elabscience Bionovation Inc. A Reliable Research Partner in Life Science and Medicine

Background

CD106 is a 110 kD glycosylphosphatidylinositol (GPI)-linked transmembrane protein, also known as VCAM-1 and INCAM-110. It is constitutively expressed on bone marrow stromal cells, myeloid progenitors, splenic dendritic cells, activated endothelial cells, as well as some lymphocytes. CD106 expression can be upregulated on endothelial cells by inflammatory cytokines. CD106 is involved in adhesion and acts as a counter-receptor for VLA-4 ( $\alpha$ 4/ $\beta$ 1 integrin) and LPAM-1 ( $\alpha$ 4/ $\beta$ 7 integrin).

For Research Use Only