

A Reliable Research Partner in Life Science and Medicine

# Recombinant FDPS/Farnesyl Diphosphate Synthase Monoclonal Antibody

catalog number: AN300076P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity Human

Immunogen Recombinant Human FDPS / Farnesyl Diphosphate Synthase protein

HostRabbitIsotypeIgGCloneA1186PurificationProtein A


**Buffer** 0.2 μm filtered solution in PBS

### Applications Recommended Dilution


WB 1:500-1:2000 FCM 1:25-1:100 ICC/IF 1:20-1:100

**IP** 1-4 μL/mg of lysate

#### Data





Flow cytometric analysis of Human FDPS expression on HeLa cells. The cells were stained with purified anti-Human FDPS, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.



Immunofluorescence analysis of Human FDPS in HUVEC cells. Cells were fixed with 4% PFA, permeabilzed with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human FDPS Monoclonal Antibody (1:60) at 37°C 1 hour. Then cells were stained with the Alexa Fluor® 488-conjugated goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI for nuclear staining (blue). Positive staining was localized to cytoplasm.

Rev. V1.2





 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

## **Elabscience®**

### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Immunoprecipitation analysis using 2  $\mu$ L anti-FDPS Monoclonal Antibody and 15  $\mu$ l of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate using FDPS Monoclonal Antibody at a dilution of 1:100. Lane A:0.5 mg Hela Whole Cell Lysate, Lane B:0.5 mg HepG2 Whole Cell Lysate, Lane C:0.5 mg A549 Whole Cell

Lysate

Western Blot with FDPS / Farnesyl Diphosphate Synthase Monoclonal Antibody at dilution of 1:500. Lane A: HepG2 Whole Cell Lysate, Lane B: Hela Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW:38 kDa Calculated-MW:48 kDa

Observed-MW:38 kDa Calculated-MW:48 kDa

### **Preparation & Storage**

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when

stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

### **Background**

This gene encodes an enzyme that catalyzes the production of geranyl pyrophosphate and farnesyl pyrophosphate from isopentenyl pyrophosphate and dimethylallyl pyrophosphate. The resulting product, farnesyl pyrophosphate, is a key intermediate in cholesterol and sterol biosynthesis, a substrate for protein farnesylation and geranylgeranylation, and a ligand or agonist for certain hormone receptors and growth receptors. Drugs that inhibit this enzyme prevent the post-translational modifications of small GTPases and have been used to treat diseases related to bone resorption. Multiple pseudogenes have been found on chromosomes 1, 7, 14, 15, 18 and X. Multiple transcript variants encoding different isoforms have been found for this gene.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.2