

## Elab Fluor® 647 Anti-Mouse CD123 Antibody[5B11]

Catalog Number: AN00962M

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2a
<b>Clone No.</b>	5B11
<b>Isotype Control</b>	Elab Fluor® 647 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832M]
<b>Conjugation</b>	Elab Fluor® 647
<b>Conjugation Information</b>	Elab Fluor® 647 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

**FCM** Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



Staining of C57BL/6 murine bone marrow cells with FITC

Anti-Mouse/Human CD11b Antibody[M1/70] and Elab Fluor® 647 Anti-Mouse CD123 Antibody[5B11] (left) or Elab Fluor® 647 Rat IgG2a, κ Isotype Control (right). Total viable cells were used for analysis.

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	IL-3 Receptor α chain;IL-3Rα
<b>Uniprot ID</b>	Q8CII2
<b>Gene ID</b>	16188

### For Research Use Only

## Background

CD123 is a 70 kD  $\alpha$  chain subunit of the IL-3 receptor (IL-3R  $\alpha$ ). It is a member of the immunoglobulin superfamily that is expressed on hematopoietic progenitors, basophils, mast cells, and megakaryocytes. This transmembrane glycoprotein can bind IL-3 with low affinity but cannot transduce signals without association with additional protein partners. CD123 can complex with either the common  $\beta$  chain (CDw131) or the IL-3R  $\beta$  chain (AIC2A) to form high-affinity heterodimeric IL-3 receptors. CDw131 can complex with the  $\alpha$  subunits of the mouse IL-3R, IL-5R and GM-CSFR to form high-affinity receptors, while the IL-3 R  $\beta$  subunit is specific for IL-3 but binds with low affinity. IL-3 binding to the receptor complex can induce proliferation and differentiation of hematopoietic cells. The 5B11 antibody does not block binding of IL-3 to the high affinity IL-3 receptor.