

A Reliable Research Partner in Life Science and Medicine

# Recombinant Carbonic Anhydrase VIII/CA8 Monoclonal Antibody

catalog number: AN300188P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## **Description**

Reactivity Human

Immunogen Recombinant Human Carbonic Anhydrase VIII / CA8 protein

Host Rabbit
Isotype IgG
Clone B106
Purification Protein A


Buffer 0.2 µm filtered solution in PBS

## Applications Recommended Dilution

WB 1:500-1:1000


**IP**0.2-1 μL/mg of lysate

#### Data



Immunoprecipitation analysis using 0.5 µL anti-CA8 Monoclonal Antibody and 15 µl of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate using CA8 Monoclonal Antibody at a dilution of 1:150. Lane A:0.5 mg Hela Whole Cell Lysate, Lane B:0.5 mg A549

Whole Cell Lysate
Observed-MW:37 kDa
Calculated-MW:37 kDa



Western Blot with Carbonic Anhydrase VIII / CA8 Monoclonal Antibody at dilution of 1:500. Lane A: A549 Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW:37 kDa Calculated-MW:37 kDa

### **Preparation & Storage**

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

## **Background**

The carbonic anhydrases (or carbonate dehydratases) are classified as metalloenzyme for its zinc ion prosthetic group and form a family of enzymes that catalyze the rapid interconversion of carbon dioxide and water to bicarbonate and protons, a reversible reaction that takes part in maintaining acid-base balance in blood and other tissues. The carbonic anhydrasekl (CA) family consists of at least 11 enzymatically active members and a few inactive homologous proteins. Carbonic anhydrase protein (CA) VIII, which is a member of the CA gene family, has been shown to have no catalytic CA activity and its biological function is still unknown. Increased expression of CA-RP VIII was observed in 78% of colorectal carcinomas. It suggested that CA-RP VIII plays a role in the process of invasion in colorectal cancer.

## For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.2