



A Reliable Research Partner in Life Science and Medicine

PE/Cyanine 5.5 Anti-Human CD11c Antibody [BU15]

Catalog Number: E-AB-F1118I

Note: Centrifuge before opening to ensure complete recovery of vial contents.

_		4.0	
	escri	MTI.	on.
ш	COUL		UII

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ

Clone No. BU15

Isotype Control PE/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F097921]

Conjugation PE/Cyanine 5.5

Conjugation Information PE/Cyanine5.5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 690 nm

(e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD11 antigen-like family member C;CD11c;Integrin alpha-X;Itgax;Leukocyte adhesion

receptor p150+95

 Uniprot ID
 P20702

 Gene ID
 3687

Background CD11c is a 145-150 kD type I transmembrane glycoprotein also known as integrin αX

and CR4. CD11c non-covalently associates with integrin β2 (CD18) and is expressed on monocytes/macrophages, dendritic cells, granulocytes, NK cells, and subsets of T and B cells. CD11c has been reported to play a role in adhesion and CTL killing

through its interactions with fibrinogen, CD54, and iC3b.

Web: www.elabscience.cn