Elabscience®

Recombinant PPIL2 Monoclonal Antibody

catalog number: AN300130P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Reactivity	Human	
Immunogen	Recombinant Human PPIL2 Protein	
Host	Rabbit	
Isotype	IgG	
Clone	5D14	
Purification	Protein A	
Buffer	0.2 μm filtered solution in PBS	
Applications	Recommended Dilution	
ICC/IF	1:20-1:100	
Data		
Immunofluorescence analysis of PPIL2 in U2OS cells. Cells		
were fixed with 4% PFA, permeabilzed with 0.1% Triton X-		
100 in PBS, blocked with 10% serum, and incubated with		
rabbit anti-human PPIL2 Monoclonal Antibody (dilution ratio		
1:60) at 4°C overnight.	Then cells were stained with the	

Alexa Fluor®594-conjugated Goat Anti-rabbit IgG secondary antibody (red) and counterstained with DAPI for nuclear staining (blue).Positive staining was localized to Nucleus.

Preparation & Storage	
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag
Background	
PPIL2, is an enzyme that be	elongs to the cyclophilin family. The cyclophilins are peptidylprolyl isomerases and are highly

PPIL2, is an enzyme that belongs to the cyclophilin family. The cyclophilins are peptidylprolyl isomerases and are highly conserved ubiquitous. They play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. PPIL2 interacts with the proteinase inhibitor eglin c and is localized in the nucleus. It contains 1 PPIL2 cyclophilin-type domain and 1 U-box domain. PPIL2 accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.

Web:www.elabscience.com