CD86 Polyclonal Antibody

catalog number: E-AB-18593



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Fusion protein of human CD86

Host Rabbit
Isotype IgG


Purification Antigen affinity purification

Conjugation Unconjugated


buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1.25_1.100

Data



Western blot analysis of Human lymph gland tissue and Jurkat cell lysates using CD86 Polyclonal Antibody at dilution of 1:450



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using CD86 Polyclonal Antibody at dilution of 1:30(×200)

Observed-MV:Refer to figures Calculated-MV:38 kDa

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms.

For Research Use Only