Recombinant Human TIGIT Protein (Fc Tag)

Catalog Number: PKSH033510



	pening to ensure complete recovery of vial contents.	
Description		
Species	Human	
Mol_Mass	39.7 kDa	
Accession	Q495A1	
Bio-activity	Not validated for activity	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		

Note: Centrifuge before opening to ensure complete recovery of vial contents

kDa	MK	R
170 130 95 72 55 43 34 26		_
17	-	
10	-	Sanda Sanda

> 95 % as determined by reducing SDS-PAGE.

Background

T cell immunoreceptor with Ig and ITIM domains (TIGIT) is a member of the CD28 family within the Ig superfamily of proteins. TIGIT is expressed on NK cells and subsets of activated; memory and regulatory T cells; and particularly on follicular helper T cells within secondary lymphoid organs. It binds to CD155 and Nectin-2 that appear on dendritic cells (DC) and endothelium. Ligation of TIGIT on T cells down-regulates TCR-mediated activation and subsequent proliferation; while NK cell TIGIT ligation blocks NK cell cytotoxicity. Through CD155 and Nectin-2; which also interact with DNAM-1/CD226 and CD96/Tactile; TIGIT is part of an interacting network of Ig superfamily members that may augment or oppose each other. In particular; TIGIT binding to CD155 can antagonize the effects of DNAM1.

For Research Use Only