## SYT3 Polyclonal Antibody

catalog number: E-AB-14372




Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |  |
|--------------|--|
| Reactivity   | Human;Mouse;Rat  |
| Immunogen    | Recombinant protein of human SYT3  |
| Host         | Rabbit   |
| Isotype      | IgG  |
| Purification | Affinity purification  |
| Conjugation  | Unconjugated   |
| buffer       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

| Applications | Recommended Dilution |
|--------------|----------------------|
| IHC          | 1:50-1:200           |

Data



Immunohistochemistry of paraffin-embedded Human brain

tissue using SYT3 Polyclonal Antibody at dilution 1:50

| Preparation & Storage |   |
|-----------------------|---|
| Storage               | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.                 |
| Shipping              | The product is shipped with ice pack, upon receipt, store it immediately at the |
|                       | temperature recommended.  |

## Background

Synaptotagmins are a large gene family of synaptic vesicle type III integral membrane proteins that function as regulators of both exocytosis and endocytosis and are involved in neurotransmitter secretion from small secretory vesicles. Calcium binds to Synaptotagmin I which triggers neurotransmitter release at the synapse. Synaptotagmin II is phosphorylated by WNK1 in a process that regulates calcium-dependent interactions. Synaptotagmin IV is expressed in calcium-dependent exocytosis of secretory vesicles in endocrine cells and neurons. Synaptotagmin IV is expressed in neuronal tissues, and has the highest mRNA levels in the hippocampus. The proximity of the Synaptotagmin IV gene to markers of several psychiatric disorders suggest an involvement of synaptotagmin IV in human disease. Synaptotagmin V is a dense-core vesicle-specific protein that regulates a specific type of calcium-regulated secretion. Synaptotagmin VI interacts with adaptor protein-2 in a calcium-independent manner. Synaptotagmin VII is widely expressed in non-neuronal tissues.

## For Research Use Only