Recombinant Human IGFBP-4/IGFBP4 Protein (His Tag)

Catalog Number: PKSH032597



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| L. Company | |
|------------|----------|
| Species | Human |
| Mol_Mass | 27.0 kDa |
| Accession | P22692 |

Bio-activity Not validated for activity

| - | | | | |
|----|----|----|-----|----|
| 12 | rn | pe | 741 | AC |
| | LU | | | |

Description

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Insulin-like growth factor binding protein 4 (IGFBP-4) is a 24 kDa protein that binds insulin-like growth factor 1 (IGF-1) and IGF-2 with high affinity and inhibits IGF action in vitro. All members of the IGFBP family can bind IGF-I and IGF-II with about equal affinity, but IGFBP-4 binds IGF2 more than IGF1. It contains IGFBP N-terminal domain and thyroglobulin type-1 domain. IGFBP-4 is induced by forskolin and N6, O2'dibutyryl sdenosine 3', or 5'-cyclic monophosphate. The IGF-binding proteins can prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors.

For Research Use Only