
Recombinant XPNPEP2 Monoclonal Antibody

catalog number: AN300084P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|-----------------------------------|
| Reactivity | Human |
| Immunogen | Recombinant Human XPNPEP2 protein |
| Host | Rabbit |
| Is otype | IgG |
| Clone | 12G13 |
| Purification | Protein A |
| Buffer | 0.2 µm filtered solution in PBS |
| Applications | Recommended Dilution |
| ІНС-Р | 1:50-1:200 |
| Data | |



Immunohistochemistry of paraffin-embedded human kidney using XPNPEP2 Monoclonal Antibody at dilution of 1:100.

| Preparation & Storage | |
|-----------------------|--|
| Storage | This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles. |
| Shipping | Ice bag |
| | |

Background

Aminopeptidase P (APP) is a hydrolase specific for N-terminal imido bonds, which are common to several collagen degradation products, neuropeptides, vasoactive peptides, and cytokines. A membrane-bound and soluble form of this enzyme (XPNPEP2) have been identified as products of two separate genes. XPNPEP2, the X-linked gene that encodes membranous aminopeptidase P (APP), has been reported to associate with APP activity. The membrane aminopeptidase P (XPNPEP2) is largely limited in distribution to endothelia and brush border epithelia. APP and XPNPEP2 contain homologous blocks of sequence common to members of the "pita bread-fold" protein family, of which Escherichia coli methionine aminopeptidase is the prototype. The C-2399A variant in XPNPEP2 is associated with reduced APP activity and a higher incidence of AE-ACEi. XPNPEP2 mRNA was detected in fibroblasts that carry the translocation, suggesting that this gene at least partially escapes X inactivation. XPNPEP2 is a candidate gene for premature ovarian failure (POF).