

A Reliable Research Partner in Life Science and Medicine

### **Recombinant Cyclin E1 Monoclonal Antibody**

catalog number: AN301091L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human; Mouse; Rat

Immunogen Recombinant Human Cyclin E1 protein


HostRabbitIsotype $IgG,\kappa$ CloneB846PurificationProtein A

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.


#### Applications Recommended Dilution

**IHC** 1:400-1000 **WB** 1:1000-5000

#### Data




Western Blot with Recombinant Cyclin E1 Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: Hela cells.



Immunohistochemistry of paraffin-embedded human colon tissue using Recombinant Cyclin E1 Monoclonal Antibody at dilution of 1:200.

## Observed-MW:47 kDa Calculated-MW:47 kDa



Immunohistochemistry of paraffin-embedded rat colon tissue using Recombinant Cyclin E1 Monoclonal Antibody at dilution of 1:200.

#### **Preparation & Storage**

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

#### **Background**

#### For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Rev. V1.1

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.1