

Recombinant Cyclin E1 Monoclonal Antibody

catalog number: AN301091L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Mouse;Rat
Immunogen	Recombinant Human Cyclin E1 protein
Host	Rabbit
Isotype	Rabbit IgG
Clone	C110715
Purification	Protein A
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications

Recommended Dilution

IHC	1:400-1000
WB	1:1000-5000
IF	1:200-1000
ELISA	1:5000-20000
IP	1:50-200


Data




Western Blot with Recombinant Cyclin E1 Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: HeLa cells.

Observed-MW:47 kDa

Calculated-MW:47 kDa



Immunohistochemistry of paraffin-embedded human colon tissue using Recombinant Cyclin E1 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded rat colon tissue using Recombinant Cyclin E1 Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.4

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis.