

# Recombinant Mouse MIF Protein (His Tag)

Catalog Number:PKSM041107



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

|                                    |  |
|------------------------------------|--|
| <b>Synonyms</b>                    | Macrophage migration inhibitory factor;Delayed early response protein 6;DER6;Glycosylation-inhibiting factor;GIF;L-dopachrome isomerase;L-dopachrome tautomerase;Phenylpyruvate tautomerase; |
| <b>Species</b>                     | Mouse  |
| <b>Expression Host</b>             | E.coli   |
| <b>Sequence</b>                    | Pro2-Ala115  |
| <b>Accession</b>                   | P34884   |
| <b>Calculated Molecular Weight</b> | 13.3 kDa   |
| <b>Observed molecular weight</b>   | 11-17 kDa  |
| <b>Tag</b>                         | C-His  |

## Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 98 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | < 0.1 EU per µg of the protein as determined by the LAL method.   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from sterile PBS, pH 7.4.<br>Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual.          |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

## Data



> 98 % as determined by reducing SDS-PAGE.

## Background

Macrophage migration inhibitory factor(MIF) is a secreted protein and belongs to the MIF family. MIF is an important regulator of innate immunity. The circulating MIF binds to CD74 on other immune cells to trigger an acute immune response. Hence MIF is classified as an inflammatory cytokine. Furthermore glucocorticoids also stimulate white blood cells to release MIF and hence MIF partially counter acts the inhibitory effects that glucocorticoids have on the immune system. Finally trauma activates the anterior pituitary gland to release MIF.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017