Recombinant Human CCL17/TARC Protein (His Tag)

Catalog Number: PKSH033735



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| - | esc | | | | | . • | | |
|-----|-----|----|---|-----|---|-----|---|---|
| - 1 | Δ | Q. | | 141 | n | ш | n | m |
| ய | U | 0 | U | ш | v | UΙ | v | ш |

 Species
 Human

 Mol_Mass
 9.1 kDa

 Accession
 O92583

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

C-C motif chemokine 17 (CCL17) is a novel CC chemokine, it belongs to the intercrine beta (chemokine CC) family. CCL17 is expressed at high levels in thymus, and at a lower level in lung, colon, and small intestine. CCL17 is also transiently expressed in stimulated peripheral blood mononuclear cells. Among CC chemokine family members, CCL17 has approximately 24 - 29% amino acid sequence identity with RANTES, MIP-1 alpha, MIP-1 beta, MCP-1, MCP-2 and MCP-3. CCL17 has been identified to be Chemotactic factor for T-lymphocytes but not monocytes or granulocytes. CCL17 plays a role in T-cell development in thymus and in trafficking and activation of mature T-cells.

For Research Use Only