

Recombinant S100A7 Monoclonal Antibody

catalog number: AN300382P

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Immunogen	Recombinant Human S100A7 protein
Host	Rabbit
Isotype	IgG
Clone	11G5
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS


Applications Recommended Dilution

ICC/IF	1:20-1:100
FCM	1:25-1:100

Data



Immunofluorescence analysis of Human S100A7 in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 1% Triton X-100 in PBS, blocked with 10% serum, and incubated with Rabbit anti-Human S100A7 monoclonal antibody (1:60). Then cells were stained with the Alexa Fluor® 594-conjugated Goat Anti-rabbit IgG secondary antibody. Positive staining was localized to cytoplasm.



Flow cytometric analysis of Human S100A7 expression on HeLa cells. The cells were treated according to manufacturer's manual, stained with purified anti-Human S100A7, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein differs from the other S100 proteins of known structure in its lack of calcium binding ability in one EF-hand at the N-terminus. The protein is overexpressed in hyperproliferative skin diseases, exhibits antimicrobial activities against bacteria and induces immunomodulatory activities.

For Research Use Only