

APC Anti-Mouse NKG2A/C/E Antibody[20d5]

Catalog Number: AN00409E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Host	Rat
Isotype	Rat IgG2a, κ
Clone No.	20d5
Isotype Control	APC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	--

Data



Staining of C57BL/6 murine splenocytes cells with FITC Anti-Mouse NK1.1 Antibody and APC Anti-Mouse NKG2A/C/E Antibody[20d5] (left) or APC Rat IgG2a, κ Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	KLRC1;Killer Cell Lectin Like Receptor C1;KLRC2;Killer Cell Lectin Like Receptor C2; KLRC3;Killer Cell Lectin Like Receptor C3
Uniprot ID	P26715

For Research Use Only

Gene ID

500338

Background

The NKG2 molecules are a family of lectin-like receptors that form heterodimers with CD94. NKG2/CD94 heterodimer are primarily expressed on NK cells, and a subset of CD8+ T cells. Studies of CD94/NKG2 heterodimers on NK cells have demonstrated that the NKG2 components transduce signals after ligand binding. NKG2A transduces inhibitory signals, while NKG2C and NKG2E transduce activating signals.

For Research Use Only