Elabscience®

HIST1H2BK Polyclonal Antibody

catalog number: E-AB-52805

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Reactivity	Human; Mouse	
Immunogen	Fusion protein of human HIST1H2BK	
Host	Rabbit	
Isotype	IgG	
Purification	Antigen affinity purification	
Conjugation	Unconjugated	
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilution	
IHC	1:40-1:200	
Data		
120		and the stand of the
cancer tissue using HIS	f paraffin-embedded Human thyroid ST1H2BK Polyclonal Antibody at on of 1:60(×200)	Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using HIST1H2BK Polyclonal Antibody at dilution of 1:60(×200)
cancer tissue using HIS	ST1H2BK Polyclonal Antibody at	cervical cancer tissue using HIST1H2BK Polyclonal
cancer tissue using HIS diluti	ST1H2BK Polyclonal Antibody at on of 1:60(×200)	cervical cancer tissue using HIST1H2BK Polyclonal

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-dependent histone that is a member of the histone H2B family. The protein encoded is an antimicrobial protein with antibacterial and antifungal activity. Two transcripts that encode the same protein have been identified for this gene, which is found in the histone microcluster on chromosome 6p21.33.