

## Elab Fluor® 647 Anti-Mouse CD226 Antibody[480.1]

Catalog Number: E-AB-F1419M

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

|                                |  |
|--------------------------------|--|
| <b>Reactivity</b>              | Mouse  |
| <b>Host</b>                    | Rat  |
| <b>Isotype</b>                 | Rat IgG2a, κ   |
| <b>Clone No.</b>               | 480.1  |
| <b>Isotype Control</b>         | Elab Fluor® 647 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832M]   |
| <b>Conjugation</b>             | Elab Fluor® 647  |
| <b>Conjugation Information</b> | Elab Fluor® 647 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter). |
| <b>Storage Buffer</b>          | Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.   |


### Applications

### Recommended usage

#### FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



Staining of C57BL/6 murine splenocytes cells with FITC Anti-Mouse CD8a Antibody and Elab Fluor® 647 Anti-Mouse CD226 Antibody[480.1] (left) or Elab Fluor® 647 Rat IgG2a, κ Isotype Control (right). Total viable cells were used for analysis.

### Preparation & Storage

|                 |   |
|-----------------|---|
| <b>Storage</b>  | Keep as concentrated solution.<br>This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| <b>Shipping</b> | Ice bag   |

### Antigen Information

|                   |        |
|-------------------|--------|
| <b>Uniprot ID</b> | Q8K4F0 |
| <b>Gene ID</b>    | 225825 |

### For Research Use Only

## Background

CD226 (DNAM-1) is constitutively expressed on native CD8+ cells and on some CD4+ T cells, macrophages and NK cells. CD226 is involved in NK and T cell mediated cytotoxicity against certain tumors. CD155 and CD112 are the ligands for CD226. This antibody (480.1) is reported by the developer to partially block the binding of mouse CD155.